

Matroids and greedy algorithms

Recall minimum spanning trees and Kruskal's algorithm.

Let $G = (V, E, w)$ where $w: E \rightarrow \mathbb{R}_0$ ($w(e) \geq 0 \forall e \in E$)
be given and let $W = \max\{w(e) \mid e \in E\}$

If we set $w'(e) = W - w(e)$ then for any spanning
tree T of G we have ($n = |V|$)

$$w'(T) = (n-1)W - w(T)$$

Hence T is a minimum spanning tree wrt w



T is a maximum weight spanning tree wrt w'

From now on we look only at the maximization
version (maximum weight spanning trees)

Greedy algorithm for MST

$F \leftarrow \emptyset$

while $E \neq \emptyset$

Let $uv \in E$ have maximum weight

$E \leftarrow E - uv$

If u and v are not connected via F

$F \leftarrow F + uv$

end


Theorem The greedy algorithm finds a maximum weight spanning tree in $G=(V,E,w)$

Proof: let T_g be the greedy spanning tree and suppose $w(T_g) < w(T_{opt})$ when T_{opt} is a max weight spT of G .

let e_1, e_2, \dots, e_{n-1} be the edges that we add when T_g is constructed (in that order).

Choose T^* max weight spT and j s.t $\{e_1, e_2, \dots, e_j\} \subseteq E(T^*)$ and no max weight (*) spT contains all the edges $\{e_1, \dots, e_{j+1}\}$

$T^* + e_{j+1}$ contains a cycle C



At least one edge of C is not in T_g as this is a tree
let e be such an edge.

Then $e \in E - \{e_1, e_2, \dots, e_j, e_{j+1}\}$ so $w(e_{j+1}) \geq w(e)$
since the greedy algorithm chose e_{j+1} in step $j+1$

Now $w(T^*) \geq w(T^* - e + e_{j+1}) \geq w(T^*)$ so

$T' = T^* - e + e_{j+1}$ is a max weight spT which violates (*)

□.

Definition Let S be a finite set and

$\mathcal{F} \subseteq 2^S$ a collection of subsets of S . Then the pair $M = (S, \mathcal{F})$ is a **matroid** if the following holds

- (1) $\emptyset \in \mathcal{F}$
- (2) if $X \in \mathcal{F}$ and $Y \subseteq X$ then $Y \in \mathcal{F}$
- (3) if $X, Y \in \mathcal{F}$ and $|Y| = |X| + 1$ then $\exists y \in Y - X$ s.t. $X + y \in \mathcal{F}$

(3) says that \mathcal{F} is **hereditary**
A set Z s.t. $Z \notin \mathcal{F}$ is **dependent**
A set X s.t. $X \in \mathcal{F}$ is **independent**

Examples of Matroids

(A) $G = (V, E)$ a graph $S = E$
 $\mathcal{F} = \{E' \subseteq E \mid E' \text{ induces a forest in } G\}$

This is called the **cyclic matroid** of G

(B) S finite subset of a vector space V
say that $\{x_1, x_2, \dots, x_n\} \in \mathcal{F}$ iff the
vectors x_1, x_2, \dots, x_n are linearly independent

Property (3) of Matroids easily implies the following property which is called the **exchange property**

(3') if $X, Y \in \mathcal{I}$ and

$|Y| = |X| + k$ for some $k \geq 1$ then there are elements $y_1, y_2, \dots, y_k \in Y - X$ such that

$$X + y_1 \in \mathcal{I}, X + y_1 + y_2 \in \mathcal{I}, \dots, X + y_1 + \dots + y_k \in \mathcal{I}$$


A **base** of a matroid $M = (S, \mathcal{I})$ is a maximal independent set B , that is $B \in \mathcal{I}$ but $B + w \notin \mathcal{I} \forall w \in S - B$

Corollary

All the bases of a matroid have the same size

How do we check whether $X \in \mathcal{I}$?

We assume $M = (S, \mathcal{I})$ is given as S together with an **oracle** A



Circuits of matroids

A **circuit** of a matroid $M = (S, \mathcal{F})$ is a minimally dependent subset of S :
 Z is a circuit of $M \Leftrightarrow Z \notin \mathcal{F}$ and $Z - a \in \mathcal{F} \forall a \in Z$

Proposition let $M = (S, \mathcal{F})$ be a matroid
if $X \in \mathcal{F}$ but $X + y \notin \mathcal{F}$ then there is a
unique circuit C contained in $X + y$

proof:


$X + y$ is dependent so it contains at least one circuit

$$\text{let } C = \{a \in X + y \mid X + y - a \in \mathcal{F}\}$$

claim: C is a circuit of M

proof: As $C - a \subseteq X + y - a \in \mathcal{F} \forall a \in C$
implying that $C - a \in \mathcal{F}$ we just need to
show that $C \notin \mathcal{F}$.

suppose $C \in \mathcal{F}$. Then by (3') we can add
elements from $X - C$ to C until we have a
set $Y \in \mathcal{F}$ with $Y \in \mathcal{F}$ and $|Y| = |X|$. But
then $Y = X + y - d$ for some d and $C \subseteq Y$
implying that $d \in C$ }




We proved that C is a circuit so it only remains to prove that C is the unique circuit in $X+y$.

Suppose $D \subseteq X+y$ is another circuit

Then the picture is

X




By minimality of circuits $C \not\subseteq D$
So $\exists z \in C - D$ (and $\exists w \in D - C$)

By definition of C the set
 $X+y-z \in \mathcal{F}$ but

$D \subseteq X+y-z$ contradicts (2)
in the definition of matroids \square .

The greedy algorithm for matroids

Below we assume that the matroid $M=(S, \mathcal{I})$ is given in form of an oracle A such that



Greedy algorithm:

input: a matroid $M=(S, \mathcal{I})$ and $w: S \rightarrow \mathbb{R}_{\geq 0}$
output: a maximum weight basis of M

$B \leftarrow \emptyset, S' \leftarrow S$

while $S' \neq \emptyset$ do

let $x \in S'$ have $w(x) \geq w(y) \forall y \in S'$

$S' \leftarrow S' - x$

if $B+x \in \mathcal{I}$ then $B \leftarrow B+x$

end
output B

Theorem The greedy algorithm outputs a maximum weight basis of M .

Proof Let B_g denote the output and let B^* be an optimum basis ($w(B^*) \geq w(B) \forall \text{ basis } B$)

Let e_1, e_2, \dots, e_k be the elements added to B by the algorithm in that order.


Let $j \in [k]$ be maximum s.t. $\{e_1, e_2, \dots, e_j\} \subseteq B^*$ for some optimum basis.

If $j = k$ we are done (B_g is optimal) so assume $j < k$

Then $B^* + e_{j+1}$ contains a circuit C .

$C \not\subseteq B_g$ as C is dependent so $\exists e \in B^* - B_g$ s.t. $e \in C \cap B^*$

By the proposition, C is the unique circuit in $B^* + e_{j+1}$ so $\hat{B} = B^* + e_{j+1} - e$ is a basis



B^* is optimal so $w(B^*) \geq w(\hat{B}) \Rightarrow w(e) \geq w(e_{j+1})$ (a)

Recall that $\{e_1, e_2, \dots, e_j\} \subseteq B_g \cap B^*$ and $A = \{e_1, e_2, \dots, e_j, e\} \subseteq B^*$ so A is independent and hence e is a possible choice for the greedy algorithm in step $j+1$.

This implies that $w(e_{j+1}) \geq w(e)$ so by (a) we have $w(e) = w(e_{j+1})$ and $w(\hat{B}) = w(B^*)$, contradicting the choice of B^* as

$\{e_1, e_2, \dots, e_{j+1}\} \subseteq \hat{B}$

□.

Alternative definition of a matroid (Def 12.2 in PS)

Definition The subset system $M = (E, \mathcal{F})$ is a matroid

\Updownarrow def

The greedy algorithm finds an optimum basis for every non-negative weight function $w: E \rightarrow \mathbb{R}_{\geq 0}$

Theorem 12.5 in PS Let $M = (E, \mathcal{F})$ be a subset system

(1) M is a matroid

\Updownarrow

(2) $\forall I_p, I_{p+1} \in \mathcal{F}; |I_p| = p = |I_{p+1}| - 1$
 $\exists e \in I_{p+1} - I_p; I_{p+1} - e \in \mathcal{F}$

\Updownarrow


(3) If $A \subseteq E$ and I, I' are maximal independent subsets of A then $|I| = |I'|$

proof

(1) \Rightarrow (2):

suppose $I_p, I_{p+1} \in \mathcal{F}$; $|I_p| = p = |I_{p+1}| - 1$
but $\nexists e \in I_{p+1} - I_p$: $I_{p+1} \in \mathcal{F}$ I_p

let $w(e) = \begin{cases} p+2 & \text{if } e \in I_p \\ p+1 & \text{if } e \in I_{p+1} - I_p \\ 0 & \text{otherwise} \end{cases}$



Then $w(I_p) = p \cdot (p+2) < (p+1)^2 = w(I_{p+1})$

but G.A chooses I_p plus some 0 weight element \rightarrow that G.A. works for w .

(2) \Rightarrow (3) suppose I, I' are maximal independent subsets of A but $|I'| > |I|$

Then let $I'' \subseteq I'$ have $|I''| = |I| + 1$

and apply (2) to get that I is not a maximal subset of A } $\}$

(3) \Rightarrow (1): Suppose (3) holds but then is some non-negative w such that the G.A. does not find an optimal bn.

Let $I = \{e_1, e_2, \dots, e_i\}$ be the greedy solution where $w(e_1) \geq w(e_2) \geq \dots \geq w(e_i)$ and

$J = \{e'_1, e'_2, \dots, e'_j\}$ be an optimal solution where $w(e'_1) \geq w(e'_2) \geq \dots \geq w(e'_j)$.

Then $w(J) > w(I)$ by the assumption above

I is maximal by construction (via G.A.) and we can assume J is also maximal (by adding zero or more weight zero elements)

By (3) we have $|I| = |J|$

It suffices to prove the following claim

Claim $w(e_m) \geq w(e'_m)$ for all $m = 1, 2, \dots, i$

Proof of claim: $w(e_1) \geq w(e'_1)$ as G.A. chose e_1
suppose $\exists m \leq i$ s.t. $w(e_j) \geq w(e'_j)$ for $j = 1, 2, \dots, m-1$
but $w(e_m) < w(e'_m)$

Let $A = \{e \in E \mid w(e) \geq w(e'_m)\}$ then

$Z = \{e_1, e_2, \dots, e_{m-1}\} \subseteq A$ and Z is a maximal subnt of A (as G.A. chose $e_m \notin A$)

But $Z' = \{e'_1, e'_2, \dots, e'_m\} \subseteq A$ so it is contained in some maximal subnt Z'' of A

However now $|Z| < |Z''| \rightarrow \Leftarrow (3)$

□.

Further definitions and remarks

Let $M = (E, \mathcal{F})$ be a matroid

The **rank** of $A \subseteq E$, denoted $r(A)$ is $r(A) = \max \{|X| \mid X \subseteq A \text{ and } X \in \mathcal{F}\}$

The rank of M is the size of a base of M (recall that they all have the same size)

The set of bases of M is

$$\mathcal{B}(M) = \{B \mid B \text{ is a base of } M\}$$

Note that $\mathcal{F} = \{X \mid X \subseteq B \text{ for some } B \in \mathcal{B}(M)\}$

So $M = (E, \mathcal{F})$ is uniquely determined by its set of bases.